Repérage patients vulnérables et fragiles

*Pr Pierre Soubeyran
Institut Bergonié*

Nous avons un problème à résoudre...

...avec les patients âgés atteints de cancer

- ✦ **Mauvais pronostic**
- ✦ **Risque majoré de toxicités**
- ✦ **Toxicités inattendues**
 - Liées aux comorbidités existantes
- ✦ **Conséquences inattendues de ces toxicités**

Evaluation gériatrique

- Identification du groupe des patients vulnérables à l'aide d'une batterie de questionnaires et à l'expérience des gériatres

QUESTIONNAIRE

1425 pts	Anormal	n	%
ADL	<6	217	15.2
IADL	<8	682	47.9
GDS15	≥6	457	32.1
MMS	<24	290	20.3
MNA	≤23,5	623	43.7
Timed Get up & Go	>20 s	328	23
CIRS-G	Grade 3-4	595	41.8

Rarely

Nous avons des solutions...

364 patients	Early death < 6 m.	Functional decline	Hospitalization for toxicity
	<i>Odd ratio (95% CI)</i>	<i>Odd ratio (95% CI)</i>	<i>Odd ratio (95% CI)</i>
Events	59/339	50/299	47/354
Extension (M+ vs M-)	4.1 (1.65-10.1)		
Sex	2.62 (1.31-5.28)		
Platelet count <150 G/l			3.8 (1,3-10,8)
Clinician opinion		ns	0.51 (0,26-0,99)
PS	ns	ns	
MNA ≤ 23.5	2.91 (1.31-56.48)	ns	4.19 (1,7-10,3)
Get up and go > 20 s	2.51 (1,31-4,82)	ns	
IADL ≤ 7	ns	3 (1,13-8,09)	
GDS15 ≥ 6		2.4 (1,23-4,66)	
MMS	ns	ns	
ADL	ns		
CIRS-G			

Soubeyran JCO 2012

Hoppe JCO2013

Warkus submitted

Nous avons des solutions...

...mais elles sont disponibles pour une minorité

Evaluation gériatrique

- Identification du groupe des patients vulnérables à l'aide d'une batterie de questionnaires

Consommatrice de temps

- A peu près une heure

Disponible pour une minorité de patients

Le screening peut aider les médecins... ...à identifier les patients vulnérables

Nombreux outils

- Propriétés variables

Faisable par tous en moins de 5'

- Médecin, IDE ou ARC

Faisable dans la majorité des centres qui traitent des cancers

- Sans besoin d'expertise gériatrique

Le screening peut aider les médecins...

...à identifier les patients vulnérables

Nombreux outils

- Propriétés variables

Faisable par tous en moins de 5'

- Médecin, IDE ou ARC

Faisable dans la majorité des centres qui traitent des cancers

- Sans besoin d'expertise gériatrique

	Items	Possible answers (score)
A	Has food intake declined over the past 3 months due to loss of appetite, digestive problems, chewing or swallowing difficulties?	0 : severe decrease in food intake
		1 : moderate decrease in food intake
		2 : no decrease in food intake
B	Weight loss during the last 3 months	0 : weight loss > 3 kg
		1 : does not know
		2 : weight loss between 1 and 3 kgs
C	Mobility	0 : bed or chair bound
		1 : able to get out of bed/chair but does not go out
		2 : goes out
E	Neuropsychological problems	0 : severe dementia or depression
		1 : mild dementia or depression
		2 : no psychological problems
F	Body Mass Index (BMI (weight in kg) / (height in m ²))	0 : BMI < 19
		1 : BMI = 19 to BMI < 21
		2 : BMI = 21 to BMI < 23
H	Takes more than 3 medications per day	0 : yes
		1 : no
		0 : not as good
P	In comparison with other people of the same age, how does the patient consider his/her health status?	0.5 : does not know
		1 : as good
		2 : better
	Age	0 : >85
		1 : 80-85
		2 : <80
TOTAL SCORE		0 - 17

Le screening peut aider les médecins... ...à identifier les patients vulnérables

Table 3. Studies with a direct comparison between screening tools

First author (reference)	Number of patients	Tools	% abnormal	Sensitivity % (95% CI)	Specificity % (95% CI)	PPV % (95% CI)	NPV % (95% CI)
Soubeyran [6]	1425	G8	68	77 (74-79)	64 (59-70)	NR	NR
		VES-13	60	69 (66-71)	74 (69-79)	NR	NR
Kenis [7]	937	G8	74	87 (84-89)	59 (53-66)	86 (83-88)	61 (55-68)
		fTRST(1)	83	91 (89-93)	42 (36-48)	81 (78-84)	63 (56-71)
Biganzoli [15]	259	fTRST(2)	55	67 (64-71)	80 (74-85)	90 (87-93)	47 (42-52)
		VES-13	47	62 (54-69)	81 (71-88)	86 (79-92)	52 (44-61)
Luce [10]	211	Fried Frailty Criteria	75	87 (81-92)	49 (38-60)	77 (70-82)	66 (53-77)
		G8	80	65 (NR)	3 (NR)	44 (NR)	8 (NR)
Baitar [11]	170	ISAR	79	70 (NR)	10 (NR)	47 (NR)	22 (NR)
		KPS <80	86	29 (NR)	44 (NR)	39 (NR)	34 (NR)
Kenis [12]	140	G8	76	92 (85-96)	52 (39-65)	78 (70-84)	78 (63-88)
		GFI	47	66 (56-75)	87 (76-94)	90 (82-95)	90 (82-95)
Owusu [17]	117	G8	NR	80 (NR)	40 (NR)	83 (NR)	35 (NR)
		GFI	NR	57 (NR)	87 (NR)	94 (NR)	36 (NR)
		fTRST(1)	NR	92 (NR)	50 (NR)	87 (NR)	63 (NR)
		fTRST(2)	NR	64 (NR)	100 (NR)	100 (NR)	43 (NR)
Kellen [18]	113	VES-13	35	88 (NR)	69 (NR)	68 (NR)	88 (NR)
		KPS ≤80	38	78 (NR)	91 (NR)	87 (NR)	85 (NR)
		ECOG-PS	66	94 (NR)	55 (NR)	61 (NR)	93 (NR)
Molina-Garrido [20]	58	VES-13	49	61 (49-72)	78 (61-90)	85 (73-93)	48 (35-62)
		GFI	31	39 (28-51)	86 (70-95)	86 (70-95)	40 (29-51)
		aCGA	NR	51 (39-62)	97 (85-100)	97 (87-100)	48 (36-60)
Pottel [13]	51	VES-13	35	39 (NR)	100 (NR)	100 (NR)	18 (NR)
		Fried Frailty Criteria	35	37 (NR)	86 (NR)	95 (NR)	16 (NR)
Molina-Garrido [22]	41	G8	67	86 (72-94)	75 (52-97)	88 (NR)	71 (NR)
		VES-13	39	57 (42-71)	100 (83-100)	100 (NR)	52 (NR)
		VES-13	29	55 (NR)	100 (NR)	100 (NR)	66 (NR)
		Barber	41	59 (NR)	79 (NR)	77 (NR)	63 (NR)

Sensitivity, specificity, PPV and NPV indicate the value of the screening tool for detecting predefined abnormal geriatric assessment. CI, confidence interval; PPV, positive predictive value, NPV, negative predictive value; NR, not reported.

Items	Possible answers (score)
A Has food intake declined over the past 3 months due to loss of appetite, digestive problems, choking or swallowing difficulties?	0: severe decrease in food intake
	1: moderate decrease in food intake
	2: no decrease in food intake
B Weight loss during the last 3 months	0: weight loss > 3 kg
	1: does not know
	2: weight loss between 1 and 3 kgs
C Mobility	0: no weight loss
	1: able to get out of bed/chair but does not go out
	2: goes out
E Neuropsychological problems	0: severe dementia or depression
	1: mild dementia or depression
	2: no psychological problems
F Body Mass Index (BMI (weight in kg)/(height in m) ²)	0: BMI < 19
	1: BMI = 19 to BMI < 21
	2: BMI = 21 to BMI < 23
H Takes more than 3 medications per day	3: BMI = 23 and > 23
	0: yes
	1: no
P In comparison with other people of the same age, how does the patient consider his/her health status?	0: not as good
	1: as good
	2: better
Age	0: >85
	1: 80-85
	2: <80
TOTAL SCORE	0-17

Le screening peut aider les médecins... ...à identifier les patients vulnérables

	Items	Possible answers (score)
A	Has food intake declined over the past 3 months due to loss of appetite, digestive problems, chewing or swallowing difficulties?	0: severe decrease in food intake
		1: moderate decrease in food intake
		2: no decrease in food intake
B	Weight loss during the last 3 months	0: weight loss > 3 kg
		1: does not know
		2: weight loss between 1 and 3 kgs
		3: no weight loss
C	Mobility	0: bed or chair bound
		1: able to get out of bed/chair but does not go out
		2: goes out
E	Neuropsychological problems	0: severe dementia or depression
		1: mild dementia or depression
		2: no psychological problems
F	Body Mass Index (BMI (weight in kg) / (height in m) ²)	0: BMI < 19
		1: BMI = 19 to BMI < 21
		2: BMI = 21 to BMI < 23
		3: BMI = 23 and > 23
H	Takes more than 3 medications per day	0: yes
		1: no
P	In comparison with other people of the same age, how does the patient consider his/her health status?	0: not as good
		0.5: does not know
		1: as good
		2: better
Age	Age	0: > 85
		1: 80-85
		2: < 80
TOTAL SCORE		0 - 17

>14

32%

Bon état ou peu vulnérable

41% MGA normale
32% un Q anormal
28% > un Q anormal

≤14

68%

Probablement vulnérable

90% avec MGA anormale
Médiane: 3 questionnaires anormaux
10% avec MGA normale

Le screening peut aider les médecins... ...à identifier les patients vulnérables

>14
32%

Pas de besoin de CGA
Traitement standard

Items	Possible answers (score)
A	0: severe decrease in food intake 1: moderate decrease in food intake 2: no decrease in food intake
B	0: weight loss > 10% 1: does not lose 2: weight loss between 1 and 10% 3: no weight loss
C	0: lost or "near loss" 1: altered but still functional but does not go out 2: cannot
D	0: severe dementia or delirium 1: mild dementia or delirium 2: no psychological problem
E	0: BMI < 18 1: BMI = 18-24.9 2: BMI = 25-29.9 3: BMI = 30-34.9 4: BMI > 35
F	0: yes 1: no
G	0: not at all 1: a little 2: better
H	1: 0-10 2: 11-20 3: 21-30
TOTAL SCORE	0 - 37

≤14
68%

Nécessité d'attention spécifique
CGA à envisager
Traitement adapté à l'état du patient

Diffusion du G8 en routine quotidienne

■ *Comment rendre le questionnaire disponible?*

- Dans le dossier informatisé
- Formulaires disponibles
- Application web

■ *Qui doit le proposer au patient ?*

- Médecin
- IDE
- ' Temps d'accompagnement soignant '
- ARC

Amélioration du questionnaire

Performance d'un test **G8 modifié** avec remplacement de l'item E par l'**IADL4**

Population	Abnormal		Se	(95%CI)	Sp	(95%CI)	AUC	(95%CI)
	N (%)	Total						
Development	756 (80%)	948	77%	(74-80)	67%	(60-74)	81%	(78-84)
Internal validation	394 (81%)	486	78%	(73-82)	71%	(60-80)	83%	(79-87)
External validation	322 (95%)	338	88%	(84-91)	69%	(41-89)	90%	(84-95)

Petit A, J Ger Oncol

Intuitivement oui mais...

- Essai d'intervention nutritionnelle INOGAD
- Pas de différence malgré une intervention efficace
- Coût de mise en place
 - EGA et intervention
 - Les autorités de santé ont besoin de données

Taskforce SIOG - Méthodologie

Elaboration des questions auxquelles répondre

	Intervention	Question
1	Geriatric interventions	Does diet intervention improve nutrition parameters ?
2		Does exercise intervention improve physical functioning ?
3		Does diet and exercise intervention improve quality of life ?
4		Does exercise intervention improve other dimensions of patients' outcome ?
5		Does geriatric advices improve appropriate application of treatment plan ?
6		Does geriatric domain-specific intervention (physical, nutritional ...) improve survival ?
7		Does geriatric domain-specific intervention improve fatigue ?
8	Patient education	Does it improve quality of life ?
9		Does it improve other dimensions of patients' outcome ?
10	Case management	Does it improve survival ?
11		Does it improve quality of life ?
12		Does it improve other dimensions of patients' outcome ?

Evaluation de chaque essai

- Niveaux d'évidence ASCO pour l'objectif principal

Table 1 – ASCO levels of evidence and grades of recommendation ⁸

Level	Type of evidence
I	Evidence obtained from meta-analysis of multiple, well-designed, controlled studies or from high-power randomised, controlled clinical trial
II	Evidence obtained from at least one well-designed experimental study or low-power randomised, controlled clinical trial
III	Evidence obtained from well-designed, quasi-experimental studies such as non-randomised, controlled single-group, pre-post, cohort, time or matched case-control series
IV	Evidence obtained from well-designed, non-experimental studies such as comparative and correlational descriptive and case studies
V	Evidence obtained from case reports and clinical examples

- ✦ **Regroupement des essais par question**
- ✦ **Synthèse en grade de recommandation (ASCO)**

Table 1 – ASCO levels of evidence and grades of recommendation ^B

Grade	Type of supporting evidence
A	There is evidence of type I or consistent findings from multiple studies of type II, III, or IV
B	There is evidence of type II, III, or IV and findings are generally consistent
C	There is evidence of type II, III, or IV but findings are inconsistent
D	There is little or no systematic empirical evidence

10 essais spécifiquement chez des patients de plus de 60 ans

Study ID	First author	Journal	Year	Report	Study design	N° of patients	Population	Age	Mean age	Intervention	Control	End point	Results
A1	Katsi T	J Cancer	2015	Article	Prospective cohort comparison study	135	Older cancer patients with cancer at the start of chemotherapy	>70 yrs	78.6 yrs	Screening questionnaire to stratify patients into low or high risk CDA in high risk patients	Routine care	1-year mortality	Significant difference in completion of chemotherapy toxicity, completion of cancer treatment as planned, treatment modifications, early treatment discontinuation, dose escalations, death at 6 months (p=0.155). No effect on mortality (p=0.96).
A2	Bourdel-Marchasson I	J Geriatr Oncol / Plos One	2015	Abstract / Article	Control trial	338	Patients with solid tumour treated by chemotherapy at risk of malnutrition (<23.5)	>70 yrs	77.5 yrs	4-month diet counselling intervention	Usual care	Postoperative delirium, postoperative complications, mortality, length of hospital stay, return to preoperative living situation and care, care dependency and quality of life	No significant difference for delirium, complications, mortality, length of stay, care dependency and QoL. Significant lower bodily pain at discharge than at admission. Significant difference between groups for return to preoperative living situation and care.
A3	Hempenius I	Plos One	2013	Article	Control trial	32	Patients with a confirmed diagnosis of prostate cancer receiving ADT but no other treatment	>70 yrs	70 yrs	12-month single-blind randomized controlled trial	Usual care	Change in functional status (baseline / 12-month and 24-month) assessed using the Medical Outcomes Study Short Form 36	Significant change in functional status between intervention group and control (p=0.01): amelioration of functional decline in intervention group. Significant change in diet quality, physical activity and significant change in physical function (p=0.007) between groups.
A4a	Demark-Wahnefeld W	J Clin Oncol	2012	Article	Control trial	82	Sedentary men with a prostate cancer who received continuous ADT for a minimum of 6 months prior to recruitment and with planned long-term retention on ADT	No specific age	71 yrs	12-week combined supervised exercise (aerobic and resistance exercises) and dietary advice (macro/nutrients) component. Supervised exercise twice per week from weeks 1-6 and once per week from weeks 7-12. Self-directed independent exercise sessions at least one per week from week 1-6 and twice per week from week 7-12. Healthy eating supervised exercise program (one session per week)	Usual care	Muscle strength, objective physical function, self-reported physical function, fatigue	Significant difference between the intervention and the ITT analysis (p=0.03), leg press (per protocol analysis p=0.03), (per protocol analysis and ITT analysis by EORTC QLQ-C15 analysis only (p=0.02)). No significant difference for chair stand (p=0.07 and p=0.09), p=0.31), PFD (p=2.88 and p=0.52), LFQI overall function (p=0.38 and p=0.21), LFQI basic upper-extremity function (p=0.68 and p=0.13), LFQI advanced upper-extremity function (p=0.32 and p=0.28) and fatigue (p=0.12 and p=0.06).
A4b	Morley MC	JAMA	2010	Article	Control trial	100	Patients radically operated for lung cancer living within a radius of 80 km from the hospital	>70 yrs	64.63 yrs	12-week combined supervised exercise (aerobic and resistance exercises) and dietary advice (macro/nutrients) component. Supervised exercise twice per week from weeks 1-6 and once per week from weeks 7-12. Self-directed independent exercise sessions at least one per week from week 1-6 and twice per week from week 7-12. Healthy eating supervised exercise program (one session per week)	Usual care	Disease-specific quality of life, change in diastolic blood pressure, patient-reported fatigue, patient exercise behaviour, aerobic exercise tolerance, body mass index, dietary behaviour	Significant difference between the intervention and the control group for exercise behaviour at 12 wk (p=0.001) and 6 mo (p=0.028), disease-specific QoL at 12wk (p=0.001) and 6 mo (p=0.007), aerobic exercise tolerance at 12 wk (p=0.001) and 6 mo (p=0.001).
A5	Lapid MI	Palliat Support C	2014	Article	Control trial	78	Postmenopausal women >50 yrs with diagnosis of stage 0-3a breast cancer one year post chemotherapy/regularly participating in resistance and/or impact exercise in the past month	>62 yrs	62 yrs	12-month exercise program, resistance plus impact intervention group: two supervised classes and one home-based session per week of progressive resistance training	Usual care	Body composition, muscle strength	Significant difference at 4 months for bodily pain of SF36 (p=0.01) and at 1 year for general health (p=0.01) and vitality components of SF36. No difference at 4 months and 1 year for the 6 min walk test (4 months (p=0.17), 1 year (p=0.93)), dyspnoea (p=0.24), FEV1 (p=0.85 and p=0.80) and FVC (p=0.12).
A6	Demark-Wahnefeld W	J Clin Oncol	2014	Article	Control trial	100	Prostate cancer patients with no structured exercise	No specific age	71.7 yrs	12-month supervised resistance and aerobic training	Pedometer and modified educational booklet with general recommendation of cardiovascular fitness, physical function, health-related quality of life	Significant difference between the intervention and the control group at 6 mo and 12 mo for cardiorespiratory fitness (p=0.029 and p=0.028), physical function (p=0.006 and	
A7	Rao V	J Cancer Surv	2014	Article	Control trial	106	Prostate cancer patients with no structured exercise	No specific age	71.7 yrs	12-month supervised resistance and aerobic training	Pedometer and modified educational booklet with general recommendation of cardiovascular fitness, physical function, health-related quality of life	Significant difference between the intervention and the control group at 6 mo and 12 mo for cardiorespiratory fitness (p=0.029 and p=0.028), physical function (p=0.006 and	
A8	Dobek J	J Cancer Surv	2014	Article	Control trial	106	Prostate cancer patients with no structured exercise	No specific age	71.7 yrs	12-month supervised resistance and aerobic training	Pedometer and modified educational booklet with general recommendation of cardiovascular fitness, physical function, health-related quality of life	Significant difference between the intervention and the control group at 6 mo and 12 mo for cardiorespiratory fitness (p=0.029 and p=0.028), physical function (p=0.006 and	
A9	Gavino DA	J Cancer Surv	2014	Article	Control trial	100	Prostate cancer patients with no structured exercise	No specific age	71.7 yrs	12-month supervised resistance and aerobic training	Pedometer and modified educational booklet with general recommendation of cardiovascular fitness, physical function, health-related quality of life	Significant difference between the intervention and the control group at 6 mo and 12 mo for cardiorespiratory fitness (p=0.029 and p=0.028), physical function (p=0.006 and	

25 essais avec un âge médian de plus de 60 ans

Selon les domaines d'intervention

Interventions physiques et état physique

Intervention physique

- 15 essais (dont 2 spécifiques sujets âgés)
- Trois de niveau I
- 11 significatifs

A There is evidence of type I or consistent findings from multiple studies

Intervention gériatrique

- 2 essais (dont 1 spécifique sujets âgés)
- 2 niveaux II
- 2 négatifs

Case Management

- 4 essais spécifiques sujets âgés
- 3 niveaux II, 1 niveau III
- 1 positif, 3 négatifs

> 14 ←

Standard management

Items	Possible answers (score)
A	Has food intake declined over the past 3 months due to loss of appetite, digestive problems, chewing or swallowing difficulties?
	0: severe decrease in food intake
	1: moderate decrease in food intake
	2: no decrease in food intake
B	Weight loss during the last 3 months
	0: weight loss > 3 kg
	1: 2 kg and above
	2: weight loss between 1 and 2 kg
	3: no weight loss
C	Mobility
	0: bed or chair bound
	1: able to get out of bed/chair but does not go out
	2: goes out
	3: severe dementia or depression
E	Neuropsychological problems
	0: severe dementia or depression
	1: no psychological problems
F	Body Mass Index (BMI) (weight in kg) / (height in m) ²
	0: BMI < 19
	1: BMI = 19 to BMI < 21
	2: BMI = 21 to BMI < 23
	3: BMI = 23 and > 23
H	Takes more than 1 medication per day
	0: yes
	1: no
P	In comparison with other people of the same age, how does the patient consider his/her health status?
	0: not as good
	1: does not know
	2: good
	3: better
	4: > 85
	5: 65-84
	6: 45-64
	7: < 45
	8: > 17
	9: < 17
	10: > 17
	11: < 17
	12: > 17
	13: < 17
	14: > 17
	15: < 17
	16: > 17
	17: < 17
	18: > 17
	19: < 17
	20: > 17
	21: < 17
	22: > 17
	23: < 17
	24: > 17
	25: < 17
	26: > 17
	27: < 17
	28: > 17
	29: < 17
	30: > 17
	31: < 17
	32: > 17
	33: < 17
	34: > 17
	35: < 17
	36: > 17
	37: < 17
	38: > 17
	39: < 17
	40: > 17
	41: < 17
	42: > 17
	43: < 17
	44: > 17
	45: < 17
	46: > 17
	47: < 17
	48: > 17
	49: < 17
	50: > 17
	51: < 17
	52: > 17
	53: < 17
	54: > 17
	55: < 17
	56: > 17
	57: < 17
	58: > 17
	59: < 17
	60: > 17
	61: < 17
	62: > 17
	63: < 17
	64: > 17
	65: < 17
	66: > 17
	67: < 17
	68: > 17
	69: < 17
	70: > 17
	71: < 17
	72: > 17
	73: < 17
	74: > 17
	75: < 17
	76: > 17
	77: < 17
	78: > 17
	79: < 17
	80: > 17
	81: < 17
	82: > 17
	83: < 17
	84: > 17
	85: < 17
	86: > 17
	87: < 17
	88: > 17
	89: < 17
	90: > 17
	91: < 17
	92: > 17
	93: < 17
	94: > 17
	95: < 17
	96: > 17
	97: < 17
	98: > 17
	99: < 17
	100: > 17
	101: < 17
	102: > 17
	103: < 17
	104: > 17
	105: < 17
	106: > 17
	107: < 17
	108: > 17
	109: < 17
	110: > 17
	111: < 17
	112: > 17
	113: < 17
	114: > 17
	115: < 17
	116: > 17
	117: < 17
	118: > 17
	119: < 17
	120: > 17
	121: < 17
	122: > 17
	123: < 17
	124: > 17
	125: < 17
	126: > 17
	127: < 17
	128: > 17
	129: < 17
	130: > 17
	131: < 17
	132: > 17
	133: < 17
	134: > 17
	135: < 17
	136: > 17
	137: < 17
	138: > 17
	139: < 17
	140: > 17
	141: < 17
	142: > 17
	143: < 17
	144: > 17
	145: < 17
	146: > 17
	147: < 17
	148: > 17
	149: < 17
	150: > 17
	151: < 17
	152: > 17
	153: < 17
	154: > 17
	155: < 17
	156: > 17
	157: < 17
	158: > 17
	159: < 17
	160: > 17
	161: < 17
	162: > 17
	163: < 17
	164: > 17
	165: < 17
	166: > 17
	167: < 17
	168: > 17
	169: < 17
	170: > 17
	171: < 17
	172: > 17
	173: < 17
	174: > 17
	175: < 17
	176: > 17
	177: < 17
	178: > 17
	179: < 17
	180: > 17
	181: < 17
	182: > 17
	183: < 17
	184: > 17
	185: < 17
	186: > 17
	187: < 17
	188: > 17
	189: < 17
	190: > 17
	191: < 17
	192: > 17
	193: < 17
	194: > 17
	195: < 17
	196: > 17
	197: < 17
	198: > 17
	199: < 17
	200: > 17

1500 patients to be screened

≤ 14

Randomisation

est bienvenu pour participer

Physicians
behavior

Immunosenescence

Alternative treatments

Predictors of toxicity
and outcome

Biomarkers of frailty
and longevity

Cost effectiveness

Tumor behavior

Patients
behavior

Patients and
family education